

Mixing Instructions

With **KBC 30/35 SA & KLC 10 (code: 969452 + 969502)**

Country **Kenya**
Specie **Layers**

Batch weight **1.000 kg**

9200/9204



Version:	20-12-2021	960359 Galdus PreStarter	201-9200 Starter	202-9200 Grower	203-9204 Developer	204-9204 Pre layer	205-9204 Pre Peak	206-9204 Layer 1	207-9204 Layer 2	
Code	Ingredient	Unit (kg)	0-10 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	28-45 weeks	45 weeks - end
960359	Broiler Prestarter Galdus High Altitude	Min 100gr/DOC								
969452	KBC 35/30 SA			350	300					
969502	KLC 10					100	100	100	100	100
KN627	MAIZE 8,5%CP			520	510	565	580	580	600	600
KN699	SOYBEANMEAL 46%CP					30	60	80	85	60
KN718C	SUNFLOWERCAKE 25 CP						50	50	75	75
KN673C	WHEAT BRAN/POLLARD			125	185	290	150	100	30	50
KN768	SOYBEAN OIL			5	5		5	10	10	5
KN778C	LIMESTONES 33.5					15	55	80	100	110
	Total			1.000	1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	22,0	21,4	20,1	14,7	15,2	15,3	15,2	14,4
002	Crude Fat	%	4,6	4,5	4,5	4,1	4,8	5,1	5,3	4,8
003	Crude Fibre	%	4,6	4,1	4,5	5,3	5,5	5,1	5,1	5,2
089	Dry matter	%	88,7	88,2	88,2	88,1	88,7	89,0	89,2	89,2
004	Ash	%	5,8	6,0	5,7	5,7	9,2	11,5	13,2	14,2
098	ME Poultry	Kcal	2900	2880	2840	2760	2760	2760	2760	2690
016	Total lysin	%	1,31	1,29	1,17	0,71	0,76	0,79	0,79	0,73
017	Total Methionin	%	0,54	0,53	0,48	0,38	0,40	0,40	0,41	0,40
019	Total Methionin + Cystin	%	0,92	0,88	0,82	0,65	0,67	0,67	0,67	0,65
048	Calcium	%	0,85	0,90	0,78	0,85	2,20	3,04	3,71	4,04
049	Total Phosphor	%	0,57	0,57	0,57	0,57	0,51	0,48	0,45	0,45
145	Digestible Phosphor	%	0,46	0,42	0,38	0,34	0,33	0,32	0,31	0,31
052	Sodium	%	0,18	0,16	0,14	0,15	0,15	0,15	0,15	0,15
050	Chloride	%	0,27	0,29	0,26	0,22	0,22	0,22	0,22	0,22
208	Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209	Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC	Cocciostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

*DOC: Day Old Chick

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as cocciostat and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- In case of no oil; replace the oil with corn. Note that FCR is likely to increase.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- Make sure birds always have access to fresh, cool and clean water.
- In case of management related issues, ask for the Koudijs Management Guide.

V10_06