

Mixing Instructions

With **KBC 30/35 SA & KLC 20 CARO (code: 969452 + 969564)**
 Country **Kenya**
 Specie **Layers**
 Batch weight **1.000 kg**

9200/9202



Version: 20-12-2021	960359 Galdus PreStarter	201-9200 Starter	202-9200 Grower	203-9202 Developer	204-9202 Pre layer	205-9202 Pre Peak	206-9202 Layer 1	207-9202 Layer 2	
Code Ingredient	Unit (kg)	0-10 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	28-45 weeks	45 weeks - end
960359 Broiler Prestarter Galdus High Altitude	Min 100gr/DOC								
969452 KBC 35/30 SA			350	300					
969564 KLC 20 + Caro					180	200	200	200	200
KN627 MAIZE 8,5%CP			520	510	550	575	585	590	585
KN673C WHEAT BRAN/POLLARDS			125	185	250	160	120	100	100
KN768 SOYBEAN OIL			5	5		5	10	10	5
KN778C LIMESTONES 33.5					20	60	85	100	110
Total			1.000	1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001 Crude Protein	%	22,0	21,4	20,1	16,5	16,1	15,5	15,2	15,2
002 Crude Fat	%	4,6	4,5	4,5	4,1	4,4	4,8	4,7	4,2
003 Crude Fibre	%	4,6	4,1	4,5	4,9	4,1	3,7	3,5	3,5
089 Dry matter	%	88,7	88,2	88,2	88,1	88,4	88,6	88,7	88,7
004 Ash	%	5,8	6,0	5,7	6,0	9,6	11,8	13,2	14,1
098 ME Poultry	Kcal	2900	2880	2840	2770	2780	2780	2760	2700
016 Total lysin	%	1,31	1,29	1,17	0,78	0,79	0,77	0,76	0,76
017 Total Methionin	%	0,54	0,53	0,48	0,40	0,41	0,41	0,40	0,40
019 Total Methionin + Cystin	%	0,92	0,88	0,82	0,69	0,69	0,68	0,67	0,67
048 Calcium	%	0,85	0,90	0,78	0,94	2,30	3,13	3,63	3,97
049 Total Phosphor	%	0,57	0,57	0,57	0,55	0,50	0,47	0,46	0,46
145 Digestible Phosphor	%	0,46	0,42	0,38	0,33	0,33	0,32	0,32	0,32
052 Sodium	%	0,18	0,16	0,14	0,13	0,15	0,15	0,15	0,15
050 Chloride	%	0,27	0,29	0,26	0,25	0,27	0,27	0,27	0,27
208 Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209 Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

*DOC: Day Old Chick

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- In case of no oil; replace the oil with corn. Note that FCR is likely to increase.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- Make sure birds always have access to fresh, cool and clean water.
- In case of management related issues, ask for the Koudijs Management Guide.

V10_06