

## Mixing Instructions

With **KBC 30/35 SA & KLC 10 CARO (code: 969452 + 969502)**  
 Country **Kenya**  
 Specie **Layers**  
 Batch weight **1.000 kg**



**Economy/9201+9205**

Version: 10-3-2022	960359	201-9201	202-9201	203-9204	204-9205	205-9205	206-9205	207-9205
Code Ingredient	Galdus PreStarter HA 0-10 days	Starter 1-6 wks	Grower 7-12 wks	Developer 13-16 wks	Prelayer 17-19 wks	Pre-peak 20-28 wks	Layer 1 28-45 wks	Layer 2 45wks-End
960359 Broiler Prestarter Galdus High Altitude	min. 100gr/DOC							
969452 KBC 35/30 SA		350	300					
969502 KLC 10				100	100	100	100	100
KN627 MAIZE 8,5%CP		540	520	565	595	605	605	600
KN673C WHEAT BRAN/POLLARDS		110	180	290	150	30	20	30
KN699C SOYBEANMEAL 47.8CP				30	45	80	80	70
KN718C SUNFLOWERCAKE 25 CP					60	100	100	100
KN778C LIMESTONES 33.5 CA				15	50	85	95	100
<b>Total</b>		<b>1.000</b>	<b>1.000</b>	<b>1.000</b>	<b>1.000</b>	<b>1.000</b>	<b>1.000</b>	<b>1.000</b>

## CALCULATED ANALYSIS

001 Crude Protein	%	22,0	21,3	20,2	14,7	15,0	15,8	15,6	15,3
002 Crude Fat	%	4,6	4,0	4,0	4,2	4,5	4,7	4,6	4,6
003 Crude Fibre	%	4,6	4,0	4,4	5,4	5,7	5,8	5,7	5,7
089 Dry matter	%	88,7	88,1	88,1	88,1	88,6	89,1	89,2	89,2
004 Ash	%	5,8	6,0	5,7	5,7	8,8	11,9	12,8	13,3
098 ME Poultry	Kcal	2900	2870	2820	2760	2760	2730	2720	2690
016 Total lysin	%	1,31	1,29	1,17	0,72	0,74	0,81	0,81	0,78
017 Total Methionin	%	0,54	0,53	0,48	0,38	0,40	0,42	0,42	0,41
019 Total Methionin + Cystin	%	0,92	0,88	0,82	0,65	0,67	0,70	0,69	0,68
048 Calcium	%	0,85	0,90	0,78	0,85	2,03	3,22	3,55	3,72
049 Total Phosphor	%	0,57	0,56	0,57	0,57	0,52	0,47	0,46	0,46
145 Digestible Phosphor	%	0,46	0,42	0,38	0,34	0,33	0,32	0,32	0,32
052 Sodium	%	0,18	0,16	0,14	0,15	0,15	0,15	0,15	0,15
050 Chloride	%	0,27	0,29	0,26	0,22	0,22	0,22	0,22	0,22
208 Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209 Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

\*DOC: Day Old Chick

### Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- When oil is available; replace 0,5% corn for 0,5% oil.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- Make sure birds always have access to fresh, cool and clean water.
- In case of management related issues, ask for the Koudijs Management Guide.