

Mixing Instructions

With **KBC 30/35 SA & KLC 20 CARO (code: 969452 + 969564)**
 Country **Kenya**
 Specie **Layers**
 Batch weight **1.000 kg**



Economy/ 9201 + 9203

Version: 3-1-2023	960359 Galdus PreStarter	201-9201 Starter	202-9201 Grower	203-9202 Developer	204-9203 Pre layer	206-9203 Layer 1	207-9203 Layer 2		
Code	Ingredient	Unit (kg)	0-10 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-45 weeks	>45 weeks
960359	Galdus High Altitude	Min. 100 gr/DOC							
969452	KBC 35/30 SA			350	300				
969564	KLC 20 + Caro					180	200	200	200
KN627	CORN 8,5%CP			540	520	550	590	630	610
KN673C	WHEAT BRAN/POLLARDS			110	180	250	150	70	80
KN778C	LIMESTONES 33.5					20	60	100	110
	Total			1.000	1.000	1.000	1.000	1.000	1.000

CALCULATED ANALYSIS

001	Crude Protein	%	22,0	21,3	20,2	16,5	16,0	15,1	15,1
002	Crude Fat	%	4,6	4,0	4,0	4,1	3,9	3,8	3,7
003	Crude Fibre	%	4,6	4,0	4,5	4,9	4,0	3,3	3,4
089	Dry matter	%	88,7	87,8	87,9	88,1	88,3	88,5	88,7
004	Ash	%	5,8	6,1	5,7	6,0	9,6	13,1	14,1
098	ME Poultry	Kcal	2900	2870	2820	2770	2760	2750	2700
016	Total lysin	%	1,31	1,29	1,17	0,78	0,79	0,75	0,75
017	Total Methionin	%	0,54	0,53	0,48	0,40	0,41	0,40	0,40
019	Total Methionin + Cystin	%	0,92	0,88	0,82	0,69	0,69	0,67	0,67
048	Calcium	%	0,85	0,93	0,80	0,94	2,30	3,63	3,97
049	Total Phosphor	%	0,57	0,53	0,55	0,55	0,50	0,44	0,45
145	Digestible Phosphor	%	0,46	0,42	0,38	0,33	0,33	0,32	0,32
052	Sodium	%	0,18	0,16	0,14	0,14	0,15	0,15	0,15
050	Chloride	%	0,27	0,29	0,26	0,26	0,27	0,27	0,27
208	Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added
209	Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added

*DOC: Day Old Chick

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 100 gram/bird total, preferably more).
- When oil is available; replace 0,5% corn for 0,5% oil.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- Make sure birds always have access to fresh, cool and clean water.
- In case of management related issues, ask for the Koudijs Management Guide.

V10_06