

OPTIMAL MIXING INSTRUCTIONS LAYER CONCENTRATES KLC 10

For feed delivering the best technical performance for the highest profits

A CONTRACTOR OF THE STATE OF	Pre -starter	Starter	Glowei	Developei	Fie Layer	FIG-FGAK	Layer	Layer 2
Ingredients	0 - 10 days	1 - 6 weeks	7 - 12 weeks	13 - 16 weeks	17 - 19 weeks	20 - 28 weeks	28 - 45 weeks	> 45 weeks - END
Broiler Prestarter Galdus High Altitude	Min. 100gr/DOC							
KBC 35/30 SA		350	300					
KLC 10				100	100	100	100	100
Maize 8,5% CP		520	510	565	580	580	600	600
SOYBEANMEAL 46% CP				30	60	80	85	60
SUNFLOWERCAKE 25 CP					50	50	75	75
WHEAT BRAIN / POLLARD		125	185	290	150	100	30	50
SOYBEAN OIL		5	5		5	10	10	5
LIMESTONES 33.5				15	55	80	100	110
Total Kgs		1000	1000	1000	1000	1000	1000	1000

IMPORTANT REMARKS:

- DOC stands for Day Old Chick
- For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 150 gram/bird total, preferably more).
- When oil is available; replace 0,5% corn for 0,5% oil.
- Always make sure all birds have access to enough feed and water.
- Carefully follow the provided mixing instructions and thoroughly mix the ingredients for the best results.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the birds houses only.
- Ensure proper quality of the limestone is used. Quicktests are available, ask your local representative for more info.
- To avoid drop in feed intake; ensure the change of feed (from Starter to Grower feed etc.) happens gradually.
- Do not modify the instructions without consulting a De Heus advisor.
- Please check the website for the full nutritional composition of the feed after mixing.

SCAN THE QR CODES TO LEARN MORE ABOUT OUR PRODUCTS











